

# The Use of Antidiabetic Herbal Remedies by Jordanian Herbalist: A Comparison of Folkloric Practice vs. Evidence-Based Pharmacology

Reem Issa<sup>1\*</sup>, Areen Khattabi<sup>2</sup>, Tabarak A. Alkarem<sup>3</sup>, Omar Altameemi<sup>3</sup>

<sup>1</sup> Department of Pharmaceutical Chemistry and Pharmacognosy.

<sup>2</sup> Department of Pharmaceutics and Pharmaceutical Sciences.

<sup>3</sup> Undergraduate Students .Faculty of Pharmacy.

## ABSTRACT

Traditionally used herbal remedies are an essential part of the Jordanian folklore, due to its abundance in vast sources of ethnomedicine.

This study conducted a face to face interview with a random sample of local herbalists, focused on traditionally used antidiabetic herbal remedies. Subsequently, these remedies were validated for evidence-based pharmacological studies, using data in the literature to find out the available information about different clinical, *in vitro* or *in vivo* studies.

Two traditionally used antidiabetic herbal mixtures were found commonly prescribed, composed of twelve different plant species. Nine species proved to be effective in *in vitro*, *in vivo* and clinical evaluations for their antidiabetic effect. Only one species was found to be effective on animal models or as enzyme inhibitor, but not in any clinical studies. Another two species were found to be ineffective neither in *in vitro*, *in vivo* nor in clinical evaluations.

This review will help to strengthen the relation between traditional medicine and evidence-based pharmacology. Also, it will shed the light on the ethnomedicine practices in Jordan, revealing some common errors among herbalist's practices and uses of antidiabetic herbal remedies. We recommend for these traditionally used plants, to go for extensive chemical investigations as well as clinical trials, in order to find out potential new antidiabetic agents.

**Keywords:** Antidiabetic, Ethnomedicine, Evidence-based, Jordan, Pharmacology.

## 1. INTRODUCTION

Traditional uses of medicinal herbal remedies have been of interest not only in Jordan, but also at the Arab world and at the global level.<sup>1</sup> Historically, a lot of ethnobotanical surveys revealed that medicinal plants are used to treat varied diseases; like the nervous system, cardiovascular systems, respiratory system, digestive and other systems.<sup>2</sup> Because of being natural, there are common believes that they are highly effective and safe.<sup>3</sup> Nevertheless, many previous as well as recent research and reviews revealed that there is a significant gap exists between scientific validation of the traditional uses, and the proved pharmacological activities of herbs used by the

public and herbalists.<sup>(2,4,5,6)</sup>

As the alarming rising trend of diabetes prevalence in the Arab region constitutes a real challenge for health decision makers,<sup>(7)</sup> a similar scenario was also shown in Jordan. In a study performed by Al-Nsour and coworkers in 2012 on Jordanian population with chronic diseases, it was estimated that 11% of the study population in 2007 was diagnosed with diabetes by a health professional, and 19% was diagnosed with diabetes according to laboratory testing.<sup>(8)</sup> In addition; diabetes were also found to cause the proportion of deaths attributable to diabetes in Jordan to increase up to 7% in 2010.<sup>(9)</sup> Therefore, more research is needed to further discover new potential antidiabetic agents. One of the possible approaches is via searching for remedies that are commonly used among the local herbalists and diabetic patients in the country.<sup>(10)</sup>

The Mediterranean location of Jordan makes it one of

\* r\_issa@asu.edu.jo

Received on 6/5/2018 and Accepted for Publication on 4/12/2018.

the richest countries in the world with a diverse floral source, which can be used in different aspects of medicine.<sup>(11)</sup> In addition, cultural beliefs and geographical diversity of the country also encourage the treatment of various diseases with plants in different formulations like crude extracts, whole plants, a paste of plants, infusions, etc.<sup>(12,13)</sup> In an ethno-pharmacological survey of medicinal herbs in Jordan at the Northern Badia region by Alzweiri *et al.*; rated 10 herbal species that possess anti-diabetic effect as the third among the list of commonly used herbs, after digestive and respiratory problems.<sup>(11)</sup> Among the previous ethno-pharmacological studies performed in Jordan, 30 different plant species were used as anti-diabetic or hypoglycemic agents in the folk medicine; with only 6 species were not reported in the literature for their claimed uses.<sup>(14)</sup>

In a cross-sectional study conducted by Otoom *et al.* they interviewed diabetic patients visiting two medical centers at the north of Jordan. In that study they found that 31% of interviewed patients have used herbal products, and that the most commonly used herbs were *Trigonella foenumgraecum* (22.9%).<sup>(15)</sup> Where in a similar study by Wazaify *et al.*, interviewing diabetes patients in Amman (the capital of Jordan), only 16.6% reported using herbs and the most commonly used was green tea (20.5%).<sup>(16)</sup>

Variations among the previous studies can be explained by the differences in their geographic locations, as in the capital Amman, patients are less familiar with the traditional uses of medicinal herbs. Moreover, the ethno-pharmacological practices by the herbalists allocated in Amman, are not well scientifically validated. Therefore, in the current article, we have surveyed for traditional herbal remedies commonly prescribed by the local herbalists allocated in Amman, for their diabetic customers, with their methods of preparation and use. The aim of this work is to review different clinical, invitro or invivo studies performed on these remedies, which have been published through various reputed scientific journals; in order to validate the reported traditional uses, and hence to assess

whether or not these practices are supported by any evidence-based pharmacological studies and/ or medicinal reports.

## **Methodology**

### **Ethnobotanical survey with the local herbalists**

We conducted a random sample of herbalist from local community of Amman areas (n=15), over a period of 2 months in 2017. Interviews were face to face with the herbalists at their shops, without time limit and according to ISE code of ethics ([www.ethnobiology.net/ethics.php](http://www.ethnobiology.net/ethics.php)). Arabic language was the language used. The interviewees were aware of their right to refuse answering any question, to stop the interview at any time, or to simply decline the interview altogether.

In order to determine the traditionally used plant species with antidiabetic effect; data collection regarding the used herbal remedies was comprised of: mixture components, local Arabic common names of the plants, and the method of preparation and use. The identity of each plant species mentioned by herbalists was not able to be checked and confirmed by a botanist; as they were available only as powder materials.

The local Arabic names of the plants were used to search for the botanical names of these species, utilizing previously published studies on the available medicinal plant species in the Jordanian market as reported by the local researchers.<sup>(17, 13, 18, 11)</sup> These botanical names were then searched in the literature for evidence-based pharmacological studies and medicinal reports supporting their claimed antidiabetic effect.

### **Literature Review**

A literature review was conducted for the plant species that have been used as antidiabetic remedies by the interviewed herbalists. The available information about different invivo, invitro and clinical antidiabetic studies of these species were collected from various electronic sources like Pubmed, SciFinder, Elsevier, Springer,

Scopus, Science Direct, Google Scholar and Web of Science, apart from these locally available books and peer reviewed journal were also used to collect information. The results of this review will be used to evaluate the herbalist's level of knowledge on the herbal remedies that are commonly prescribed by them for their diabetic customers.

## Results

### Ethnobotanical survey results

As a result of our ethnobotanical survey, two mixture herbal remedies composed of 12 plant species, used for treatment of diabetes were reported. The Arabic common names, botanical names, and methods of preparation for

each mixture are shown in Table 1. These species were found to belong to 11 families. Of the reported plant families, Compositae, Brassicaceae, Fabaceae, Asteraceae, Oleaceae and Lamiaceae were previously reported to possess antidiabetic activity.<sup>(19)</sup> Moreover, only 7 plant species were previously reported as antidiabetic agents by the local researchers namely: *Oleae uropaea* and *Salvia triloba* by Alzweiri *et al.*; <sup>(11)</sup> *Artemisia vulgaris*, *Artemisia herba-alba*, *Rheum ribes*, *Teucrium polium*, and *Cinnamomum ceylanicum* by Abu-Irmaileh and Afifi.<sup>13</sup> Up to our knowledge, the other five plant species were not previously reported to be used as antidiabetic agents among the local herbalists in Jordan.

Table (1)

The most commonly used antidiabetic herbal mixtures by the local herbalists

*Plants botanical name	Family name	Local Arabic name
<b>Mixture 1:</b> Take a spoon and then boil and drink, repeated three times a day after each meal		
<i>Artemisia vulgaris L.</i>	Compositae	البعيثران
<i>FucusvesiculosusL.</i>	Fucaceae	الفوقس الحويصلي
<i>Lepidiumsativum L.</i>	Brassicaceae	الرشاد
<i>AstragalusonbrgchisL.</i>	Fabaceae	الكثيراء
<i>Artemisia herba-alba Asso</i>	Asteraceae	الشيح
<i>Oleaeuropaea L.</i>	Oleaceae	ورق الزيتون
<b>Mixture 2:</b> Take a spoon and then boil and drink on empty stomach before bed time		
<i>Artemisia vulgaris L.</i>	Compositae	البعيثران
<i>Artemisia herba-alba Asso</i>	Asteraceae	الشيح
<i>Rheum ribesL.</i>	Polygonaceae	الروباص
<i>TeucriumpoliumL.</i>	Compositae	الجعدة
<i>UrticapiluliferaL.</i>	Urticaceae	القريص
<i>Salvia triloba L.</i>	Lamiaceae	الميرمية
<i>CostusigneusNak</i>	Costaceae	القسط الهندي
<i>Cinnamomumceylanicum</i>	Lauraceae	القرفة

\*Plants botanical name corresponding to the local Arabic name as prescribed by previous researchers. <sup>(17, 13, 18, 11)</sup>

### Literature validation for the featured plant species

Studying and evaluating the local herbalist level of knowledge on using herbal remedies is considered one of

the most important issues, since modulating their practices and communicating with target group can be more effective in the treatment process. Therefore, over the past

few decades, a lot of reviews have been done investigating the traditional uses and scientific studies related to the globally or locally antidiabetic herbs used, from which we can validate the efficient use of herbal remedies by the herbalist's for our featured plants species. Those species are discussed below in term of their efficiency in controlling diabetes. Moreover, these plants phyto-constituents and mode of action for their antidiabetic effect were also reviewed.

### 1. Antidiabetic activity of *Artemisia vulgaris*

As an antidiabetic herb, *A. vulgaris* has been reported in Jordan to be commonly recommended by the local herbalists for the treatment of diabetes.<sup>(20, 13)</sup> However, it has been noticed that when mice are treated with a dosage of 5 mg/ 20 g extracts of *A. vulgaris*, there was no significant lowering in their blood glucose levels.<sup>(21)</sup>

Phytochemical analysis identified 15 out of a total of 22 compounds, all are being classified as phenolic compounds; mainly chlorogenic acid derivatives or flavonoids, with a high radical-scavenging capacity and a strong reducing potential in the methanolic extract.<sup>(22)</sup>

### 2. Antidiabetic activity of *Fucus vesiculosus*

The extracts obtained from *F. vesiculosus* algae have been traditionally used for the treatment of obesity and several gastrointestinal diseases.<sup>(23)</sup> For instance, in a study performed by Gabbia *et al.*, the ability of the extracts to inhibit the digestive enzymes (like  $\alpha$ -amylase and  $\alpha$ -glucosidase) and to control the postprandial plasma glucose levels has been evaluated in a mouse model of non-alcoholic steato-hepatitis.<sup>(24)</sup> The obtained results demonstrated that both the postprandial glycemc peak and the area under curve for the blood glucose were reduced using these algae phytocomplex extracts. This algal extract may also be useful in the control of carbohydrate digestion and absorption, since a delay in carbohydrate digestion resulted from the administrations of the extract in a diet particularly rich in fat, even though a decrease in its assimilation might also occur.

The effect of brown marine algae as hypoglycemic

agent was attributed to the presence of fucoidan, a negatively charged water-soluble polysaccharide. Depending on the algal species from which fucoidan was extracted as well as the algae's season of harvest, fucoidan inhibits  $\alpha$ -glucosidase differently.<sup>(25)</sup> Furthermore, Wang *et al.* showed that if fucoidan is orally administered for 13 weeks, a decrease in fasting blood glucose, blood urea nitrogen, serum creatinine, urine protein and collagen levels in non-obese Wistar rat will be observed.<sup>(26)</sup> These findings suggest that fucoidan has the ability to prevent or impede the development of diabetic nephropathy related to spontaneous diabetes and attenuate hyperglycemia compared with the control Wistar rats.

The effect of *F. vesiculosus* on post challenge plasma glucose and insulin levels in men and women has been investigated by Paradis *et al.* using a randomized crossover placebo-controlled trial.<sup>(27)</sup> Compared with placebo, administration of 250 mg seaweed capsules resulted in a 12.1% reduction in the insulin incremental area under the curve and a 7.9% increase in the Cederholm index of insulin sensitivity. This, indeed, indicates that insulin homeostasis may be altered by brown seaweed in response to carbohydrate ingestion.

### 3. Antidiabetic activity of *Lepidium sativum*

The hypoglycemic effect due to *L. sativum* seeds was investigated in normal and streptozotocin induced diabetic rats. A significant decrease in blood glucose levels was produced in diabetic and normal rats after an acute or chronic repeated oral administration of aqueous extract, with no changes in basal plasma insulin concentrations; meaning that this pharmacological activity is based most likely on a mechanism that is independent of insulin secretion.<sup>(28)</sup> Later, the same research group has performed a similar study and concluded that the aqueous *L. sativum* extract caused a significant inhibition of renal glucose reabsorption, which in turn reduced blood sugar.<sup>(29)</sup> Moreover, the ethanolic extracts of *L. sativum* showed a high in vitro antioxidant activity, where a higher scavenging activity was exhibited by shoots compared to

the seed. This is most likely due to the higher content of the total polyphenolic compounds in the Leaf extract.<sup>(30)</sup>

#### 4. Antidiabetic activity of *Astragalus onbrgchis*

It has been concluded that *A. onbrgchis* total saponins and curcumin, with different distribution ratio, can significantly improve lipid profile and reduce glycosylated serum protein.<sup>(31)</sup> Moreover, it increases the high density lipoprotein (HDL) content and serum insulin levels; and therefore it can be used for the prevention and treatment of diabetes. Furthermore, *Astragalus* polysaccharides was used to treat rats with type 2 diabetes mellitus, the results suggested that the expression level of miRNA-203a-3p was up regulated in their liver, indicating insulin resistance attenuation in type 2 diabetes.<sup>(32)</sup>

In a large meta-analysis study including 1054 participants, a control group was compared with *Astragalus* aqueous decoction treated group. That study found that *Astragalus* administration can significantly reduce both fasting and postprandial plasma glucose when compared with control group; thus it may be beneficial as a supplementary therapy in the treatment of type 2 diabetes.<sup>(33)</sup>

#### 5. Antidiabetic activity of *Artemisia herba alba* Asso

*Artemisia herba alba* is one of the most commonly used herbal plants to treat diabetes in Saudi Arabia.<sup>(34)</sup> In Jordan, as well, this plant has been reported as an antidiabetic herb with both in vitro and in vivo hypoglycemic activity.<sup>(35, 20, 36, 13)</sup>

When the ethanolic extract of *A. herba alba* aerial part was given orally to both normal and diabetic albino rats, the blood glucose level was significantly reduced in type II diabetic rat.<sup>(37)</sup> A similar study performed by Boudjelal *et al.* showed that when *A. herba alba* infusion were administered orally to normal and alloxan-induced diabetic rats, a significant reduction in blood glucose content with a much more efficient antidiabetic activity compared to glibenclamide was observed.<sup>(38)</sup> In addition, a recent study has shown that the aqueous extracts of *A. herba alba* exhibited an  $\alpha$ -amylase inhibitory activity.<sup>(39)</sup>

This effect was most likely related to the high phenolic and flavonoidal contents in the extract, which is composed of several polyphenol compounds; with 5-caffeoylquinic (chlorogenic) acid being the main compound.

A considerable lowering of elevated blood sugar was obtained when fifteen patients with diabetes mellitus were treated with *A. herba alba* extract; 14 out of the 15 patients had good remission of diabetic symptoms. More importantly, no side effects were reported either during or after treatment with the extract.<sup>(40)</sup>

#### 6. Antidiabetic activity of *Olea europaea*

In Jordan, olives oil and leaves were used traditionally for the treatment of diabetes.<sup>(11)</sup> Otoom *et al.* stated that 3.1% of diabetic patients in Jordan use this plant to control their diabetes conditions.<sup>(15)</sup> In addition, Oran and Al-Eisawi and Hudaib *et al.* reported the use of *O. europaea* oil and leaves in Jordan as an antidiabetic herb with both in vitro and in vivo tested hypoglycemic activity.<sup>(41, 42)</sup>

It has been found that when extracts of *O. europaea* leaves were administrated in low and high doses, an improvement in the physiological, molecular and histopathological alterations on streptozotocin induced diabetic male rats was observed.<sup>(43)</sup> In addition, a substantial reduction in the digestion and absorption of starch was previously observed in streptozotocin induced diabetic animal models receiving olive leaf extract compared to untreated intestine.<sup>(44)</sup> These results suggest that the inhibition of disaccharide digestion at the intestinal mucosa might be the reason behind the hypoglycemic mechanism.

The aqueous extract of olive tree leaves was suggested to be potential sources of natural antioxidants and a potent inhibitory of  $\alpha$ -amylase activity, due to their high flavonoidal and phenolic contents.<sup>(39)</sup> Moreover, a study performed by Sato *et al.* showed that mice fed with a high fat diet exhibited lower serum glucose and insulin levels combined with a glucose tolerance enhancement, when exposed to oleanolic acid (triterpene) from *O. europaea*.<sup>(45)</sup> In addition to this, oleic acid, phenols and squalene, were

all attributed to the use of plant extract in the treatment of diabetes, hypertension and toothache.<sup>(46, 36, 47)</sup>

Another compound that is also responsible for the biological activities of olive is oleuropeoside, which exhibited an antidiabetic activity in alloxan-induced diabetic animals; with suggested mechanisms for the hypoglycaemic activity involve an increase in peripheral uptake of glucose and also potentiating of glucose-induced insulin release.<sup>(48)</sup>

In a controlled randomized clinical trial, diabetic patients were treated orally with 500 mg of olive leaf extract tablet once daily.<sup>(44)</sup> The results showed that HbA1c and fasting plasma insulin levels were significantly lowered after 14 weeks as compared to the control group.

#### 7. Antidiabetic activity of *Rheum ribes*

*Rheum ribes* is most commonly used in the treatment of diabetes in Jordan.<sup>(13)</sup> The effect of the water extract was found to exert a dual inhibition of  $\alpha$ -amylase and  $\alpha$ -glucosidase in a significant dose dependent manner, using an in vitro enzymatic starch digestion bioassay.<sup>(49)</sup> Also, it was found that the aqueous extract of *R. ribes* to possess glucose homeostasis and diabetic neuropathy protection in alloxan-diabetic mice. Moreover high doses can significantly increase serum catalase and decrease glucose levels.<sup>(50)</sup> Interestingly, in the same study, the mice were also found to be protected from acquiring diabetes and diabetic neuropathy as they were pretreated with the extract prior to alloxination. Another study performed by Hamzeh and coworkers showed that alloxan-induced diabetic rats exhibited an improvement in their renal dysfunction; due to the effect of hydroalcoholic extract of *R. ribes* root, through controlling blood glucose and renal protective effects.<sup>(51)</sup>

#### 8. Antidiabetic activity of *Teucrium polium*

Different studies had confirmed the use of *Teucrium polium* in Jordan as anti-diabetic herb, with both in vitro and in vivo tested hypoglycemic activity.<sup>(52, 35, 41, 20, 53, 36, 13, 11)</sup>

Recently, Al-Kharabsheh *et al.* demonstrated that the

aqueous extract of the Jordanian species of *T. polium* exhibited weak antioxidant and  $\alpha$ -amylase inhibitor; due to their relatively low phenolic and flavonoidal contents.<sup>(39)</sup> These findings are consistent with the previous study by Kasabri *et al.*, who found that the water extract of *T. polium* does not have any in vitro  $\alpha$ -amylase inhibitory effect.<sup>(54)</sup>

On the contrary, in a more detailed study, the ethanol extract of this plant were found to exhibit a tendency to decrease glucose level and increase blood insulin level in streptozotocin diabetic rats.<sup>(55)</sup> It seems that, as was further confirmed by Stefkov *et al.*, the ethanolic extract has an insulinotropic effect on INS-1E cells and a reduction in blood glucose levels in both normo- and hyperglycemic rats, due to the presence of flavonoids.<sup>(56)</sup> These findings are in agreement with Mirghazanfari *et al.* and Patel *et al.*, who showed that the crude extract of *T. polium* is able to enhance insulin secretion by the pancreas, due to the presence of apigenin which exists only in methanol fraction but not in aqueous fractions.<sup>(57,58)</sup> In addition, it seems that this type of extract has an ability to regenerate the islets of Langerhans comparing to the untreated diabetic rats. From these data, we may conclude that the hypoglycemic effect of *T. polium* extract varies as it is largely influenced by the type of the solvent used for extracting the active components.

In a clinical trial, the hypoglycemic effect of this plant was compared with glybenclamide in diabetes mellitus type 2.<sup>(59)</sup> The results showed that the mean of glycosylated hemoglobin decreased in both treatments; where the mean levels of triglyceride, total cholesterol and body mass index decreased significantly in plant extract treated group, with decreased appetite. On the other hand, HDL-Cholesterol, LDL-Cholesterol and liver function tests showed no change in both groups.

#### 9. Antidiabetic activity of *Urtica pilulifera* (*Urtica dioica*)

It was reported that about 8% of diabetic patients in Jordan use *Urtica pilulifera* (Nettle) as antidiabetic herb

with in vitro/in vivo tested hypoglycemic activity and  $\alpha$ -amylase/ $\alpha$ -glucosidase inhibitory effects.<sup>(15)</sup> On the contrary, Hamdan and Afifi could not observe any  $\alpha$ -glucosidase inhibitory activity for *U. dioica* aqueous extract;<sup>(60)</sup> while Oenal *et al.* were previously able to demonstrate this inhibitory effect on  $\alpha$ -glucosidase for the same plant species.<sup>(61)</sup>

Similarly, when different extracts of *U. pilulifera* were prepared and tested for their antidiabetic, antioxidant and antiinflammatory effects on type 2 diabetic rat model, big conflicts in their results were observed, which can be explained by the type of the tested plant species that were not clearly distinguished in the literature. The extracts of *U. dioica* showed an ability to enhance insulin secretion by islets of Langerhans and reduce blood sugar levels.<sup>(62)</sup> On the other hand, no significant decrease in glucose movement in the gastrointestinal tract in an in vitro study was observed.<sup>(63)</sup> Similarly, not any hypoglycemic effect in alloxan-diabetic rats was observed; however a significant antihyperglycemic effect (lasting for 3 hours) was observed after the oral glucose tolerance test.<sup>(64)</sup> This indicates that the intestinal glucose absorption was reduced due to aqueous *U. dioica* extract effect.

While the hexan extract of *U. dioica* showed no clear effect, both ethyl acetate and chloroform extracts exhibited a significant antiinflammatory, hypoglycemic and antioxidant effects in diabetic rats.<sup>(65)</sup> Another recent study evaluated the synergistic effects of hydro-alcoholic extract of *U. dioica* combined with pioglitazone on the prevention of diabetic nephropathy in streptozotocin induced-diabetic mice.<sup>(66)</sup> The results showed that this combination have synergistic protective effects that can be a promising approach for the treatment of diabetic nephropathy.

In a randomized double-blind placebo-controlled clinical trial, a capsule of (500 mg) obtained from Nettle leaf extract and administrated daily to patients with type 2 diabetes, showed that the blood levels of fasting glucose and HbA1c were significantly reduced, without any

significant effects on the other parameters compared with placebo.<sup>(67)</sup>

#### **10. Antidiabetic activity of *Salvia triloba* (*S. fruticosa*) (*S. officinalis*)**

In a study by Otoom *et al.* they reported the use of *S. officinalis* by (3.1%) of diabetic patients in Jordan.<sup>(16)</sup> Al-Mustafa and Al-Thunibat revealed the use of *S. fruticosa* by 54% of the traditional healers in Jordan for treatment of diabetes and its complications.<sup>(68)</sup> Moreover, the study showed that the aqueous and the methanol extracts to contain total phenolic compounds of 18.5% and 46.7% respectively, with moderate antioxidant effects. Alzweiri *et al.* reported the use of *S. triloba* in Jordan as antidiabetic herb.<sup>(11)</sup>

These variations in the type of species used within the same study area were clearly reported by Kasabri *et al.*<sup>(69)</sup> Briefly, eleven Jordanian indigenous *Salvia* spp., were found to modulate gastrointestinal carbohydrate and lipid digestion and absorption, when were screened in vitro for inhibition of pancreatic triacylglycerol lipase,  $\alpha$ -amylase and  $\alpha$ -glucosidase using the crude aqueous extracts, in dose dependent manner. These data explain the appearance of more than one *Salvia* species in the literature; that have been traditionally used for treatment of diabetes either by the traditional healers or the citizens in Jordan.

In a study by Perfumi *et al.* investigating the hypoglycemic activity of leaf infusion of *S. fructose* in normo- and alloxan-diabetic rabbits, this infusion was found to cause significant reduction in blood glucose levels in diabetic rabbits without exerting any effect on normal ones.<sup>(70)</sup> Additionally, the hypoglycemic effect was dependent on the root of administration as changing it from oral to intravenous, rendered the extracts of *S. fructose* ineffective; thus indicating that this plant would reduce the intestinal absorption of glucose leading to hypoglycemic effect.

In a randomized placebo-controlled trial performed on hyperlipidemic type 2 diabetic patients, treated with 500 mg encapsulated hydroalcoholic extract of *S.*

*officinalis*, a significant reduction in the blood levels of glucose, HbA1c, total cholesterol, triglyceride, and LDL; and an increase of HDL level were reported.<sup>(71)</sup> Similarly, a randomized placebo-controlled trial performed on type 2 diabetic patients, treated with 150 mg *S. officinalis* extract, a significant reduction of 2 h postprandial glucose and total cholesterol levels were reported but with no observed effects on fasting glucose, HbA1c, triglyceride, LDL and HDL.<sup>(72)</sup>

### 11. Antidiabetic activity of *Costus igneus* (*Costus spicatus* D. Don)

In in vitro study the ethanolic extract of *C. spicatus* leaf was used to evaluate glucose uptake activity, showed no direct peripheral action comparable with insulin and metformin.<sup>(73)</sup> A similar previous study investigating insulin tolerance, showed that *C. spicatus* tea consumption did not alter insulin sensitivity in obesity induced hyperglycemic mice, which suggested that *C. spicatus* leaves is not effective in the treatment of obesity-induced hyperglycemia.<sup>(74)</sup>

Phytochemical screening of *C. igneus* and Insulin plant (*C. spicatus*) leaves revealed that it is rich in protein, iron, and antioxidant components such as ascorbic acid,  $\alpha$ -tocopherol,  $\beta$ -carotene, terpenoids, steroids, and flavonoids, where the methanolic extract varied in its content of phytochemicals including carbohydrates, triterpenoids, proteins, alkaloids, tannins, saponins, and flavonoids.<sup>(75)</sup> Moreover, the insulin-like protein (ILP) was purified from *C. igneus* grown in Western Ghats of India by affinity chromatography using anti-insulin antibodies. The characterization of ILP showed that it is structurally different from insulin but functionally similar. Interestingly, the purified ILP showed significant hypoglycemic effect; when administered orally in oral glucose tolerance test, compared to insulin in streptozotocin induced diabetic mice.<sup>(76)</sup>

In a cross-sectional clinical study, patients consuming either one fresh leaf or 1 teaspoon of shade-dried powder/day of *C. igneus* in conjunction with other

modalities of treatment had effectively produced glycemic control in diabetics.<sup>(77)</sup>

### 12. Antidiabetic activity of *Cinnamomum ceylanicum*

Abu-Irmaileh and Afifi reported the use of *C. ceylanicum* in Jordan as antidiabetic herb.<sup>(13)</sup> In vitro incubation of pancreatic islets with cinnamaldehyde isolated from *C. ceylanicum*, resulted in enhanced insulin release. The insulinotropic effect of cinnamaldehyde was due to increase in the glucose uptake through glucose transporter translocation in peripheral tissues.<sup>(78)</sup>

A group of type 2 diabetic patients were treated with glibenclamide, and randomly assigned to receive either 1 g of cinnamon or placebo daily for 12 weeks. A highly significant reduction of fasting blood glucose level after 6 and 12 weeks of treatment (10.12% and 17.4%, respectively), was observed. Compared to baseline value and to placebo group at corresponding duration, a similar effect on glycosylated Hb was also found. These findings indicate the beneficial effect of adjuvant cinnamon as antidiabetic agent along with the conventional medications, in order to treat poorly controlled type 2 diabetes mellitus.<sup>(79)</sup>

### Discussion

In the current survey, 9 out of the 12 listed plant species were found clinically studied for their antidiabetic effects. For the rest of the species (*A. vulgaris*, *L. sativum*, *R. ribes*) only invitro and in vivo studies were available, with not any clinical studies were performed on these species.

Nine species, *F. vesiculosus*, *A. onbrgchis*, *A. herba alba*, *O.europaea*, *T. polium*, *U. Pilulifera*, *S. triloba*, *C. igneus*, and *C. ceylanicum* proved to be effective in in vivo, invitro and in clinical studies as hypoglycemic agents, or as adjuvant treatments in order to reduce diabetic complications. Of the other species, *L. sativum* was found effective in invitro/ in vivo studies, but not clinically proved for its antidiabetic effectiveness. In spite that the species *R. ribes* and *A. vulgaris*, failed to show any hypoglycemic effects in invitro or in in vivo studies; they



showed some evidences to reduce diabetes complications when tested on diabetic animal's models.

The phytoconstituents analysis data of the effective species are in agreement with the previously published data, Mishra *et al.* summarized the most naturally abundant antidiabetic phytoconstituents with their mode of activity as follow: <sup>(80)</sup> Alkaloids inhibit alpha-glucosidase and decrease glucose transport through the intestinal epithelium. Imidazoline compounds stimulate insulin secretion in a glucose-dependent manner. Polysaccharides increase the levels of serum insulin, reduce the blood glucose levels and improve tolerance of glucose. Flavonoids suppress the glucose level, reduce plasma cholesterol and triglycerides significantly, and increase their hepatic glucokinase activity; probably by enhancing the insulin release from pancreatic islets. Dietary fibers effectively adsorb glucose, retard glucose diffusion, inhibit the activity of alpha-amylase, and may be responsible for decreasing the rate of glucose absorption and concentration of postprandial serum glucose. Saponin stimulates the release of insulin and blocks the formation of glucose in the blood stream. Ferulic acid has stimulatory effects on insulin secretion.

### **Significant findings**

The majority of the experiments confirmed the benefits of the listed plant species; with hypoglycaemic effects, in the management of diabetes mellitus. In addition, numerous mechanisms of actions have also been proposed for these plant species. Therefore, it is assumed that these botanicals have a major role to play in the management of diabetes, but still needs further exploration for development of drugs and nutraceuticals from natural resources.

Many herbal remedies used by the local herbalists, were found to not undergo careful scientific assessment yet. Further scientific validation of several Jordanian plant species is therefore necessarily needed, if the efficacy and safety of the botanicals in reducing diabetes level in the country are required.

### **Limitations**

The local common names of the plant species are often misleading; as there are not any exclusive authorities in the country, identifying the herbal materials commonly found at the herbalist's shops. Also, many medicinal herbs were cultivated using different breeding procedures, with the aim of increasing the production rate. Presumably, the new cultivated and hybridized crops of the same species may not correspond to the plants described in the literature. In addition, the sources of these plants are not documented to be either imported or locally harvested species, that may bring significant variations in their chemicals composition and therefore might affect their pharmacological activities.

### **Conclusion**

The current data gathered through our literature review and ethnobotanical survey; clearly showed that local herbalists have an important knowledge about the use of medicinal plants in the treatment of diabetes, as been previously proved by evidence-based pharmacological studies including clinical, invivo or invitro evaluations on some of the above mentioned species. These species could be a real natural reservoir of new biomolecule with a potential antidiabetic activity. Therefore, preservation and further investigation of this heritage by documentation and scientific evaluation, are essential requirement and should be considered as a national priority. Such findings support the necessity of proper handling of herbal medicine, which requires proper regulation and licensing as per the WHO regulations. <sup>(81, 82, 83)</sup>

This review may also help the scientists and researchers working in the field of complementary and alternative medicine, to find out a possible cure for diabetes and explore new agents with hypoglycemic effect. In addition, it may also discover false herbal treatments that have serious harmful and adverse health effects, which are caused by herb-drugs interaction or improper dosage as being described by unqualified healers, without any medical supervisions being followed on these chronically

ill patients.

### **Conflict of interest statement**

The authors declare they have no conflict of interest.

### **Acknowledgments**

We would like to give thanks to Applied Science

Private University, Amman, Jordan; for the continuous support. Also, the authors thank all respondents for their important contribution at this survey by sharing their knowledge on the use of medicinal plants.

## **REFERENCES**

- (1) . Azaizeh H, Saad B, Cooper E, Said O. Traditional Arabic And Islamic Medicine, A Re-Emerging Health Aid. Evidence-Based Complementary and Alternative Medicine. 2010; 7(4):419-24.
- (2) Bone K, And Mills S. Principles and Practice Of Phytotherapy-E-Book: Modern Herbal Medicine. Elsevier Health Sciences; 2013 Jan 8.
- (3) Mukherjee PK, Bahadur S, Chaudhary SK, Kar A, Mukherjee K. Quality Related Safety Issue-Evidence-Based Validation Of Herbal Medicine Farm To Pharma. In Evidence-Based Validation Of Herbal Medicine. 2015 (Pp. 1-28).
- (4) Clement YN, Williams AF, Khan K, Bernard T, Bhola S, Fortuné M, Medupe O, Nagee K, Seaforth CE. A Gap Between Acceptance And Knowledge Of Herbal Remedies By Physicians: The Need For Educational Intervention. BMC Complementary and Alternative Medicine. 2005; 5(1):20.
- (5) Mukherjee PK, Editor. Evidence-Based Validation of Herbal Medicine. Elsevier; 2015 Feb 17.
- (6) Issa R, And Bashiti I. Herbal Medicine Use By People in Jordan: Exploring Beliefs and Knowledge of Herbalists and Their Customers. J.Biol.Sci. 2017; 17:400-409.
- (7) Boutayeb A, Lamlili ME, Boutayeb W, Maamri A, Ziyat A, Ramdani N. The Rise of Diabetes Prevalence in The Arab Region. Open Journal of Epidemiology. 2012; 2(02):55.
- (8) Al-Nsour M, Zindah M, Belbeisi A, Hadaddin R, Brown DW, Walke H. Prevalence Of Selected Chronic, Noncommunicable Disease Risk Factors In Jordan: Results of The 2007 Jordan Behavioral Risk Factor Surveillance Survey. Preventing Chronic Disease. 2012; 9.
- (9) Jarab AS, Alqudah SG, Mukattash TL, Shattat G, Al-Qirim T. Randomized Controlled Trial Of Clinical Pharmacy Management Of Patients With Type 2 Diabetes In An Outpatient Diabetes Clinic In Jordan. Journal of Managed Care Pharmacy. 2012; 18(7):516-26.
- (10) Afifi FU, Kasabri V. Pharmacological and Phytochemical Appraisal of Selected Medicinal Plants from Jordan With Claimed Antidiabetic Activities. Scientia pharmaceutica. 2013; 81(4):889-932.
- (11) Alzweiri M, Al Sarhan A, Mansi K, Hudaib M, Aburjai T. Ethnopharmacological Survey of Medicinal Herbs In Jordan, The Northern Badia Region. Journal of Ethnopharmacology. 2011; 137(1):27-35.
- (12) Lev E, Amar Z. Ethnopharmacological Survey of Traditional Drugs Sold In The Kingdom Of Jordan. Journal of Ethnopharmacology. 2002; 82(2-3):131-45.
- (13) Abu-Irmaileh BE, Afifi FU. Herbal Medicine in Jordan with Special Emphasis on Commonly Used Herbs. Journal of Ethnopharmacology. 2003; 89(2-3):193-7.
- (14) Afifi-Yazar, Fatma U.; Kasabri, Violet; Abu-Dahab, Rana. Medicinal Plants from Jordan in the Treatment of Diabetes: Traditional Uses Vs. In Vitro and In Vivo Evaluations–Part 2. Planta Medica. 2011; 77(11):1210-1220.
- (15) Ootom SA, Al-Safi SA, Kerem ZK, Alkofahi A. The Use of Medicinal Herbs by Diabetic Jordanian Patients. J Herb Pharmacother. 2006; 6: 31–41.
- (16) Wazaify M, Afifi FU, El-Khateeb M, Ajlouni K. Complementary and Alternative Medicine Use among

- Jordanian Patients with Diabetes. Complementary Therapies in Clinical Practice. 2011; 17(2):71-5.
- (17) Al-Eisawi, D.M. Vegetation of Jordan. UNESCO—Regional Office for Science and Technology for The Arab States, Cairo, 1996; 266.
- (18) Afifi, F.U., Abu-Irmaileh, B. Herbal Medicine in Jordan with Special Emphasis on Less Commonly Used Medicinal Herbs. Journal of Ethnopharmacology. 2000; 72:101–110.
- (19) Jarald E, Joshi SB, Jain DC. Diabetes vs Herbal Medicines. Iranian Journal of Pharmacology and Therapeutics. 2008; 7(1):97.
- (20) Abu Irmalah, B., Afifi, F. Treatment with Medicinal Plants in Jordan. Dirasat-Medical and Biological Sciences. 2000; 27:53–74.
- (21) Villasenor, I. M., & Lamadrid, M. R. A. Comparative Anti-Hyperglycemic Potentials of Medicinal Plants. Journal of Ethnopharmacology. 2006; 104(1-2):129-131.
- (22) Melguizo-Melguizo, D., Diaz-De-Cerio, E., Quirantes-Piné, R., Švarc-Gajić, J., & Segura-Carretero, A. The Potential of Artemisia Vulgaris Leaves as a Source of Antioxidant Phenolic Compounds. Journal of Functional Foods. 2014; 10:192-200.
- (23) Pangestuti R, Kim SK. Biological Activities and Health Benefit Effects of Natural Pigments Derived From Marine Algae. Journal of Functional Foods. 2011; 3(4):255-66.
- (24) Gabbia, D., Dall'Acqua, S., Di Gangi, I. M., Bogialli, S., Caputi, V., Albertoni, L., & De Martin, S. The Phytocomplex from Fucusvesiculosus and Ascophyllumnodosum Controls Postprandial Plasma Glucose Levels: An In Vitro and In Vivo Study in a Mouse Model Of Nash. Marine Drugs. 2017; 15(2), 41.
- (25) Kim, K. T., Rioux, L. E., & Turgeon, S. L. Alpha-Amylase and Alpha-Glucosidase Inhibition is Differentially Modulated by Fucoïdan Obtained from Fucusvesiculosus and Ascophyllumnodosum. Phytochemistry. 2014; 98, 27-33.
- (26) Wang, Y., Nie, M., Lu, Y., Wang, R., Li, J., Yang, B., & Li, X. Fucoïdan Exerts Protective Effects Against Diabetic Nephropathy Related to Spontaneous Diabetes Through The NF-Kb Signaling Pathway In Vivo and In Vitro. International Journal of Molecular Medicine. 2015; 35(4), 1067-1073.
- (27) Paradis ME, Couture P, Lamarche B. A Randomised Crossover Placebo-Controlled Trial Investigating The Effect of Brown Seaweed (Ascophyllum Nodosum and Fucus Vesiculosus) on Postchallenge Plasma Glucose and Insulin Levels in Men and Women. Applied Physiology, Nutrition, and Metabolism. 2011; 36(6):913-9.
- (28) Eddouks, M., Maghrani, M., Zeggwagh, N. A., & Michel, J. B. Study of The Hypoglycaemic Activity of Lepidium sativum L. Aqueous Extract in Normal and Diabetic Rats. Journal of Ethnopharmacology. 2005; 97(2): 391-395.
- (29) Eddouks M, Maghrani M. Effect of Lepidium Sativum L. on Renal Glucose Reabsorption and Urinary TGF-B1 Levels in Diabetic Rats. Phytotherapy Research. 2008; 22(1):1-5.
- (30) Malar, J., Chairman, K., Singh, A. R., Vanmathi, J. S., Balasubramanian, A., & Vasanthi, K. Antioxidative Activity of Different Parts of The Plant Lepidium sativum Linn. Biotechnology Reports. 2014; 3: 95-98.
- (31) Yan, X., Li, X., & Miao, M. The Intervention Effect of Different Distribution Ratio of Astragalus Total Saponins And Curcumin on The DM Rats Model. Saudi Pharmaceutical Journal. 2017; 25(4): 504-508.
- (32) Wei, Z., Weng, S., Wang, L., & Mao, Z. Mechanism of Astragalus Polysaccharides in Attenuating Insulin Resistance in Rats With Type 2 Diabetes Mellitus Via The Regulation of Liver Microrna 203a 3p. Molecular Medicine Reports. 2018; 17(1): 1617-1624.
- (33) Tian H, Lu J, He H, Zhang L, Dong Y, Yao H, Feng W, Wang S. The Effect of Astragalus as an Adjuvant Treatment in Type 2 Diabetes Mellitus: A (Preliminary) Meta-Analysis. Journal of Ethnopharmacology. 2016; 191: 206-15.
- (34) Al-Rowais NA. Herbal Medicine in the Treatment of Diabetes Mellitus. Saudi Med J. 2002; 23: 1327-1331.
- (35) Al-Khalil, S.M. A Survey of Plants Used In Jordanian Traditional Medicine. International Journal of Pharmacogenomics. 1995; 33:317–323.
- (36) Aburjai, T., Hudaib, M., Tayyema, R., Yousef, M., Qishawi, M. Ethnopharmacological Survey of Medicinal Herbs in Jordan, The Ajloun Heights Region. J.

- Ethnopharmacol. 2007; 110: 294-304.
- (37) Jemaa H.B., Khlifis., Hmed H.B., Karmous I., Benzarti A., Elati J., Et Al. Antidiabetic, Antihyperlipidemic And Antioxidant Effects Of Artemisia Herba Alba Extract On Experimental Diabetes. *Int J Pharmacol.* 2015; 11: 552-560.
- (38) Boudjelal A, Siracusa L, Henchiri C, Sarri M, Abderrahim B, Baali F, Ruberto G. Antidiabetic Effects of Aqueous Infusions of Artemisia Herba-Alba and Ajugaiava in Alloxan-Induced Diabetic Rats. *Plantamedica.* 2015; 81(09):696-704.
- (39) Al-Kharabsheh S, Al-Dabbas M, Ghazzawi H, Zatimeh A, Abulaila K. Antioxidant Activity And A-Amylase Inhibitory Effect Of Selected Medicinal Plants Grown In Jordan: An In-Vitro Study. *Journal of the Arab Society for Medical Research.* 2017; 12(1):19.
- (40) Al-Waili MSD. Treatment of Diabetes Mellitus by Artemisia Herba-Alba Extract: Preliminary Study. *Clinexp pharmacophys.* 1986; 13: 569– 573
- (41) Oran, S.A., Al-Eisawi, D.M. Check List of Medicinal Plants in Jordan. *Dirasat/Medicine and Biology Science.* 1998; 25: 84–112.
- (42) Hudaib M, Mohammad M, Bustanji Y, Tayyeb R, Yousef M, Abuirjeie M, Aburjai T. Ethnopharmacological Survey of Medicinal Plants in Jordan, Mujib Nature Reserve and Surrounding Area. *J Ethnopharmacol.* 2008; 120: 63–71.
- (43) Al-Attar, A. M., & Alsalmi, F. A. Effect of Oleae uropaea Leaves Extract on Streptozotocin Induced Diabetes in Male Albino Rats. *Saudi Journal of Biological Sciences.* 2017; in press.
- (44) Wainstein, J., Ganz, T., Boaz, M., Bar Dayan, Y., Dolev, E., Kerem, Z., & Madar, Z. Olive Leaf Extract as a Hypoglycemic Agent in Both Human Diabetic Subjects and In Rats. *Journal of Medicinal Food.* 2012; 15(7), 605-610.
- (45) Sato, H., Genet, C., Strehle, A., Thomas, C., Lobstein, A., Wagner, A., & Saladin, R. Anti-Hyperglycemic Activity of a TGR5 Agonist Isolated from Oleae uropaea. *Biochemical and Biophysical Research Communications,* 2007; 362(4): 793-798.
- (46) Silva, S., Gomes, L., Leitão, F., Coelho, A.V., Vilas Boas, L. Phenolic Compounds and Antioxidant Activity of Olea Europaea L. Fruits and Leaves. *Food Sci. Tech. Int.* 2006; 12(5), 385-396.
- (47) Waterman, E., Pharm M., Lockwood, B., Pharm, B., Phd, Pharm, S. M. R. Active Components and Clinical Applications of Olive Oil. *Altern. Med. Rev.* 2007; 12 (4): 331-342.
- (48) Gonzalez M, Zarzuelo A, Gamez MJ, Utrilla MP, Jimenez J, Osuna I. Hypoglycaemic Activity of Olive Leaf. *Planta Med.* 1992; 58: 513-515.
- (49) Kasabri V, Afifi FU, Hamdan I. In Vitro and In Vivo Acute Antihyperglycemic Effects of Five Selected Indigenous Plants from Jordan Used in Traditional Medicine. *J Ethnopharmacol.* 2011; 133:888–896.
- (50) Raafat K, Aboul-Ela M, El-Lakany A. Alloxan-Induced Diabetic Thermal Hyperalgesia, Prophylaxis and Phytotherapeutic Effects of Rheum Ribes L. in Mouse Model. *Archives of Pharmacal Research.* 2014; 26:1-0.
- (51) Hamzeh S, Farokhi F, Heydari R, Manaffar R. Renoprotective Effect of Hydroalcoholic Extract of Rheum Ribes Root in Diabetic Female Rats. *Avicenna Journal of Phytomedicine.* 2014; 4(6):392.
- (52) Dafni A, Yaniv Z, Palevitch D. Ethnobotanical Survey of Medicinal Plants in Northern Israel. *J Ethnopharmacol.* 1984; 10, 295–310.
- (53) Said O, Fulder S, Khalil K, Eli K, Bashar S. Maintaining a Physiological Blood Glucose Level With “Glucoselevel”, a Combination of Four Antidiabetes Plants Used in The Traditional Arab Herbal Medicine. *E-CAM.* 2008; 5: 421–428.
- (54) Kasabri V, Afifi FU, Hamdan I. In Vitro and in Vivo Acute Antihyperglycemic Effects of Five Selected Indigenous Plants from Jordan Used in Traditional Medicine. *J Ethnopharmacol.* 2011; 133:888–896.
- (55) Yazdanparast R, Esmaeili MA, Halen JA. Teucrium Polium Extract Effects Pancreatic Function of Streptozotocin Diabetic Rats: A Histopathological Examination. *Iranian Biomed J.* 2005; 9: 81–85.
- (56) Stefkov G, Kulevanova S, Miova B, Dinevska-Kjovkarovska S, Mølgaard P, Jäger AK, Josefsen K. Effects of Teucrium Polium Spp. Capitatum Flavonoids on The Lipid and Carbohydrate Metabolism in Rats.

- Pharmaceutical Biology. 2011; 49(9):885-92.
- (57) Mirghazanfari SM, Keshavarz M, Nabavizadeh F, Soltani N, Kamalinejad M. The Effect of “Teucrium Polium L.” Extracts on Insulin Release from in Situ Isolated Perfused Rat Pancreas in a Newly Modified Isolation Method: The Role Of Ca<sup>2+</sup> And K<sup>+</sup> Channels. *Iranian Biomedical Journal*. 2010; 14(4):178.
- (58) Patel DK, Prasad SK, Kumar R, Hemalatha S. An Overview on Antidiabetic Medicinal Plants Having Insulin Mimetic Property. *Asian Pacific Journal of Tropical Biomedicine*. 2012; 2(4):320-30.
- (59) Karimi F, Abbasi S, Bateni Ar. The Effect of Teucrium Polium on Blood Glucose in Diabetes Mellitus Type 2; A Comparison With Glibenclamide. *Iranian South Medical Journal (Ismj)*. 2002; 4 (2): 96 -103.
- (60) Hamdan I, Afifi FU. Screening of Jordanian Flora for A-Amylase Inhibitory Activity. *Pharm Biol*. 2008; 46:746–750.
- (61) Oenal S, Timur S, Okutucu B, Zihnioglu F. Inhibition of A-Glucosidase by Aqueous Extracts of Some Potent Antidiabetic Medicinal Herbs. *Prep Biochem Biotech*. 2005; 35: 29–36.
- (62) Farzami B, Ahmadvand D, Vardasbi S, Majin FJ, Khaghani S. Induction of Insulin Secretion by a Component of Urtica Dioica Leaf Extract in Perfused Islets of Langerhans and its In Vivo Effects in Normal and Streptozotocin Diabetic Rats. *J Ethnopharmacol*. 2003; 89: 47 -53.
- (63) Gallagher AM, Flatt PR, Duffy G, Abdel-Wahab YH. The Effects of Traditional Antidiabetic Plants on In Vitro Glucose Diffusion. *Nutrition Research*. 2003; 23(3):413-24.
- (64) Bnouham M, Merhfour FZ, Ziyat A, Mekhfi H, Aziz M, Leggsyer A. Antihyperglycemic Activity of The Aqueous Extract of Urtica Dioica. *Fitoterapia*. 2003; 74: 677–681.
- (65) Abo-Elmatty DM, Essawy SS, Badr JM, Sterner O. Antioxidant and Anti-Inflammatory Effects of Urtica Pilulifera Extracts in Type2 Diabetic Rats. *Journal of Ethnopharmacology*. 2013; 145(1):269-77.
- (66) Shokrzadeh M, Sadat-Hosseini S, Fallah M, Shaki F. Synergism Effects of Pioglitazone and Urtica Dioica Extract in Streptozotocin-Induced Nephropathy Via Attenuation of Oxidative Stress. *Iranian Journal of Basic Medical Sciences*. 2017; 20(5):497.
- (67) Kianbakht S, Khalighi-Sigaroodi F, Dabaghian FH. Improved Glycemic Control in Patients with Advanced Type 2 Diabetes Mellitus Taking Urtica Dioica Leaf Extract: A Randomized Double-Blind Placebo-Controlled Clinical Trial. *Clin Lab*. 2013; 59(9-10):1071-6.
- (68) Al-Mustafa AH, Al-Thunibat OY. Antioxidant Activity of Some Jordanian Medicinal Plants Used Traditionally for Treatment of Diabetes. *Pak J Biol Sci*. 2008; 11(3):351-8.
- (69) Kasabri V, Afifi FU, Abu-Dahab R, Mhaidat N, Bustanji YK, Abaza IF, Mashallah S. In Vitro Modulation of Metabolic Syndrome Enzymes and Proliferation of Obesity Related-Colorectal Cancer Cell Line Panel by Salvia Species from Jordan. *Rev Roum Chim*. 2014; 59:693-705.
- (70) Perfumi M, Arnold N, Tacconi R. Hypoglycemic Activity of Salvia Fruticosa Mill. from Cyprus. *J Ethnopharmacol*. 1991; 34:135–140.
- (71) Kianbakht S, Dabaghian FH. Improved Glycemic Control and Lipid Profile in Hyperlipidemic Type 2 Diabetic Patients Consuming Salvia Officinalis L. Leaf Extract: A Randomized Placebo. Controlled Clinical Trial. *Complementary Therapies in Medicine*. 2013; 21(5):441-6.
- (72) Behradmanesh S, Derees F, Rafieian-Kopaei M. Effect of Salvia Officinalis on Diabetic Patients. *Journal of Renal Injury Prevention*. 2013; 2(2):51.
- (73) Pareek A, Suthar M, Godavarthi A, Goyal M, Bansal V. Negative Regulation of Glucose Uptake by Costus Pictus in L6 Myotube Cell Line. *J Pharm Negative*. 2010; 1: 24–6.
- (74) Keller AC, Vandebroek I, Liu Y, Balick MJ, Kronenberg F, Kennelly EJ, Et Al. Costus Spicatus Tea Failed To Improve Diabetic Progression in C57BLKS/J Db/Db Mice, A Model Of Type 2 Diabetes Mellitus. *J Ethnopharmacol*. 2009; 121:248–54.
- (75) Hegde PK, Rao HA, Rao PN. A Review on Insulin Plant (Costus Igneus Nak). *Pharmacognosy Reviews*. 2014; 8(15):67.

- (76) Joshi BN, Munot H, Hardikar M, Kulkarni AA. Orally Active Hypoglycemic Protein from *Costus igneus* NE Br.: An in Vitro and in Vivo Study. *Biochemical and Biophysical Research Communications*. 2013; 436(2):278-82.
- (77) Shetty AJ, Parampalli SM, Bhandarkar R, Kotian S. Effect of the Insulin Plant (*Costus Igneus*) Leaves on Blood Glucose Levels in Diabetic Patients: A Cross Sectional Study. *J Clin Diagn Res*. 2010; 4: 2617–21.
- (78) Patel DK, Kumar R, Laloo D, Hemalatha S. Natural Medicines from Plant Source Used For Therapy of Diabetes Mellitus: An Overview of Its Pharmacological Aspects. *Asian Pacific Journal of Tropical Disease*. 2012; 2(3):239-50.
- (79) Sahib AS. Anti-Diabetic and Antioxidant Effect of Cinnamon in Poorly Controlled Type-2 Diabetic Iraqi Patients: A Randomized, Placebo-Controlled Clinical Trial. *Journal of Intercultural Ethnopharmacology*. 2016; 5(2):108.
- (80) Mishra SB, Rao CV, Ojha SK, Vijayakumar M, Verma A. An Analytical Review of Plants for Anti Diabetic Activity with their Phytoconstituent and Mechanism of Action. *Int J Pharm Sci Res*. 2010; 1(1):1647-52.
- (81) WHO Report on the Intercountry Expert Meeting Of Traditional Medicine and Primary Health Care, 30th November–3rd December, 1991. Cairo, Egypt.
- (82) WHO Report on the Intercountry Meeting On the Development of Guidelines for National Policy on Traditional Medicine, 21–23 December, 1993. EMRO, Alexandria, Egypt
- (83) WHO, 1999. *Monographs On Selected Medicinal Plants*, Vol. 1. WHO, Geneva.

## استخدام العلاجات العشبية المضادة للسكري من قبل العطارين الأردنيين: مقارنة بين الممارسة الشعبية والدراسات الدوائية القائمة على الأدلة

ريم عيسى<sup>1\*</sup>، عرين الخطابي<sup>2</sup>، تبارك عبد الكريم<sup>3</sup>، عمر التميمي<sup>3</sup>

<sup>1</sup> قسم الكيمياء الصيدلانية والعقاقير .

<sup>2</sup> قسم الصيدلة والعلوم الدوائية.

<sup>3</sup> طلاب جامعيين، كلية الصيدلة.

### ملخص

يعدّ التداوي بالأعشاب جزءاً أساسياً من الفلكلور الأردني وذلك بسبب كثرة استخدامه في الطب الشعبي. تم إجراء هذه الدراسة بناءً على مقابلة (وجهاً لوجه) مع عينة عشوائية من العطارين حيث تم التركيز على العلاجات العشبية المضادة لمرض السكري. وتم التحقق من صحة هذه العلاجات بالمقارنة مع الدراسات الدوائية القائمة على الأدلة العلمية، وذلك باستخدام البيانات المنشورة لمعرفة المعلومات المتاحة حول الدراسات السريرية المختلفة، والدراسات التي أجريت إما داخل الكائن الحي أو في المختبر.

ووضحت الدراسة الاستقصائية وجود اثنين من الخلطات العشبية المضادة لمرض السكري المستخدمة بشكل واسع التي تتألف من اثني عشر نوعاً من النباتات. أثبتت تسعة أنواع من هذه النباتات فعاليتها لعلاج مرض السكري كما تم تقييمها سريرياً، ومن خلال التجارب التي أجريت داخل الجسم الحي أو في المختبر. بالمقابل تم العثور على نوع واحد فقط أثبتت فعاليته على نماذج حيوانية أو كمنشط للإنزيمات، ولكن ليس في أي من الدراسات السريرية. كما تم رفض نوعين آخرين غير فعالين.

ستساعد هذه الدراسة على تعزيز العلاقة بين الطب الشعبي وعلم الأدوية القائم على الأدلة. أيضاً، سوف تلقى الضوء على ممارسات الطب الشعبي في الأردن والكشف عن بعض الأخطاء الشائعة بين العطارين في استخدام الأدوية العشبية المضادة لمرض السكري.

بناءً على ذلك فأنا نوصي ونؤكد على أهمية إجراء تجارب كيميائية وسريرية مكثفة لهذه النباتات المستخدمة في الطب الشعبي في الأردن وذلك بهدف استكشاف مركبات دوائية طبيعية لعلاج مرض السكري والتقليل من مضاعفاته.

**الكلمات الدالة:** مضادات السكري، العلاجات الشعبية، الدراسات الدوائية، الأردن.